

Quantum Hall states of atomic Bose gases: density profiles in single-layer and multi-layer geometries

N.R. Cooper

*Theory of Condensed Matter Group, Cavendish Laboratory,
Madingley Road, Cambridge CB3 0HE, United Kingdom*

F.J.M. van Lankvelt

*Rudolf Peierls Centre for Theoretical Physics,
1 Keble Road, Oxford OX1 3NP, United Kingdom*

J.W. Reijnders and K. Schoutens

*Institute for Theoretical Physics, University of Amsterdam,
Valckenierstraat 65, 1018 XE Amsterdam, the Netherlands*

(Dated: 28 July, 2005)

We describe the density profiles of confined atomic Bose gases in the high-rotation limit, in single-layer and multi-layer geometries. We show that, in a local density approximation, the density in a single layer shows a landscape of quantized steps due to the formation of incompressible liquids, which are analogous to fractional quantum Hall liquids for a two-dimensional electron gas in a strong magnetic field. In a multi-layered set-up we find different phases, depending on the strength of the inter-layer tunneling t . We discuss the situation where a vortex lattice in the three-dimensional condensate (at large tunneling) undergoes quantum melting at a critical tunneling t_{c1} . For tunneling well below t_{c1} one expects weakly coupled or isolated layers, each exhibiting a landscape of quantum Hall liquids. After expansion, this gives a radial density distribution with characteristic features (cusps) that provide experimental signatures of the quantum Hall liquids.

PACS numbers: 03.75.Lm, 73.43.Cd

I. INTRODUCTION

Among the fascinating developments in the field of quantum gases is the possibility to study correlated states of matter in a setting that is entirely different from the traditional setting of electrons in a solid state environment. A prime example are fractional quantum Hall states, which are expected when trapped atoms (bosons or fermions) are made to rotate at ultra-high angular momentum [1, 2, 3, 4, 5, 6, 7]. While there has been steady progress in achieving high angular momentum [8, 9], the conditions for the actual realization of these states have not yet been met.

The most direct experimental signature of electronic (fractional) quantum Hall states, the quantization of the Hall conductance, is not easily available for realizations of such states with neutral atoms. It is thus important to investigate the experimental signatures of atomic quantum Hall states. A number of largely complimentary proposals have been made

- fractional (braid) statistics [10]
- loss of condensate fraction [11]
- detection of gapless edge excitations [12]
- detection of density correlations in expansion image [13]
- characteristic density profiles [14, 15]

In this paper, we work out the proposals put forward in [14, 15, 16] for the detection of atomic (fractional) quantum Hall states via characteristic density profiles in a suitably engineered experimental set up.

We first analyze the case of a single layer of rotating atoms, with a weak confining potential. For this situation we find a characteristic plateau landscape for the density, with steps at sharply quantized values of the density (see Figs. 1 and 2), which are direct consequences of the incompressibility of the quantum Hall states.

After that we consider a more involved situation, where a condensate that is rotating around the z -axis is exposed to an optical lattice potential in the z -direction. This leads to a multi-layer geometry, with an inter-layer tunneling whose strength is set by the strength of the optical lattice potential. This approach is presently pursued by a number of experimental groups. With an appropriate choice of parameters, one expects a regime that can be characterized as *a stack of weakly coupled layers that each display a landscape of quantum Hall states*. We propose to probe this state after expansion in both the axial and radial directions. As the expansion of each layer in the axial direction is very fast, it gives rise to a density profile as a function of the radial distance r that is an average over the radial density profiles of the many layers. Despite the fact that the density profiles in these layers differ from each other (owing to the differing numbers of particles in the layers), we show that the resulting averaged density profile has characteristic features (cusps), which are remnants of the steps in the individual lay-

ers before expansion (see Figs. 6, 7). We propose these cusps as possible experimental fingerprints of the atomic quantum Hall states.

Our presentation is organized as follows. In section II we discuss the case of a single layer, and present the expected landscape of quantized plateaus. In section III we discuss the multi-layer set-up: section IIIB discusses the melting of the vortex lattice in the multi-layer case and section IIIC discusses the quantum phases expected after the melting. In section IIID we discuss the expansion after switching off the optical lattice and trapping potential and present the density profiles that are expected. Throughout the paper we indicate typical values of the experimental parameters that are involved, and the conditions needed to justify the approximations that we make.

II. SINGLE-LAYER GEOMETRY

A. Experimental set-up

We consider a single cloud of rotating bosons, with harmonic confinement $\omega_\perp$ in the $x - y$ (in-plane) direction and $\omega_\parallel$ in the z (out-of-plane) direction. The characteristic length scales $\ell_\perp$ and $\ell_\parallel$ are

$$\ell_{\perp,\parallel} = \sqrt{\hbar/(m\omega_{\perp,\parallel})}. \quad (1)$$

We assume that the rotation frequency ω is at or near the critical frequency $\omega_\perp$, and that parameters are such that the system is in the two-dimensional (2D) regime. In this set-up the energy scale for atom-atom interactions is

$$g_{\text{qH}} = \frac{g}{(2\pi)^{3/2}\ell_\parallel\ell_\perp^2} \quad (2)$$

with $g = \frac{4\pi\hbar^2 a_s}{m}$, where a_s is the scattering length. Assuming the values $\hbar\omega_\perp = 5\text{ nK}$, $a_s = 5\text{ nm}$, and taking $\ell_\parallel = 50\text{ nm}$ for a layer defined by an optical lattice (see below), gives an energy scale $g_{\text{qH}} \simeq 0.5\text{ nK}$. Since typically $g_{\text{qH}} < \hbar\omega_\parallel, \hbar\omega_\perp$, in the study of quantum Hall states, the single particle states can be restricted to the two-dimensional lowest Landau Level (LLL) [1]. Clearly, the observation of quantum Hall states (in single or multi-layer set-up) will require that all energy scales in the experiment (tunneling, temperature, etc.) are well below g_{qH} .

We mostly focus on one of the following two situations, which we refer to as ‘quadratic’ and ‘quartic’ in-plane confinement.

quadratic confinement. Here we assume a rotation frequency ω slightly below the critical frequency $\omega_\perp$, leaving a residual parabolic potential in the rotating frame of reference $V_2(r) = \frac{1}{2}k_2 r^2$ with $k_2 \propto (\omega_\perp - \omega)$.

quartic confinement. This is for critical rotation $\omega = \omega_\perp$, but in the presence of an additional confining potential, which we take to be a quartic, $V_4(r) = \frac{1}{4}k_4 r^4$ [8].

B. Phase separation in an external potential

We refer to Refs. [1, 2, 3, 4, 5, 6, 7] for extensive studies of the quantum liquids that form at high angular momentum, after the point where the vortex lattice melts has been crossed. Exact diagonalization studies on a disc, which represent a finite-size gas in a harmonic confinement potential (for up to $N = 10$ particles), have revealed “vortex liquid” groundstates closely related to fractional quantum Hall states[1, 2]. Due to finite size effects, the density profiles of these states show large fluctuations. Features of some (but not all) of these states have been understood in a picture invoking non-interacting composite fermions[2]. Studies in edgeless geometries (sphere or torus) have established the existence of homogeneous quantum Hall liquids[3, 4]. In these geometries the liquids have a uniform density over the entire system, so the calculations represent regions of a trapped gas where the density is approximately uniform. Studies on edgeless geometries provide the most accurate numerical data concerning the bulk properties of the quantum Hall liquids that form at high rotation rate.

In this paper we focus on inhomogeneous quantum Hall liquids, with density modulated by an external potential such as $V_2(r)$ or $V_4(r)$. The idea is the following. From the numerical studies cited earlier, we infer the existence of stable, incompressible quantum liquids at certain specific filling factors ν_i (the filling factor is defined below in terms of the ratio of the number densities of bosons and vortices), and we know the energy per particle ϵ_i in these liquids. If we now consider a given number of particles in a slowly varying external potential, we can patch together regions where the particles form specific quantum liquids, such as to minimize the total energy with the one global constraint of reproducing the total of number of particles in the trap.

For inhomogeneous systems, in which the potential $V(\mathbf{r})$ varies slowly in space compared to $\ell_\perp$, the density distribution, $n(\mathbf{r})$, (the number of atoms per unit area) averaged on scales large compared to $\ell_\perp$, can be obtained by minimizing the functional

$$E = \int d^2\mathbf{r} (e[\nu(\mathbf{r})] + V(\mathbf{r})n(\mathbf{r}) - \mu n(\mathbf{r})) , \quad (3)$$

where $e[\nu(\mathbf{r})]$ is the interaction energy per unit area, which is a function of the local filling factor $\nu(\mathbf{r}) \equiv n(\mathbf{r})/n_0$ [with $n_0 = 1/(\pi\ell_\perp^2)$], and μ is the chemical potential (which controls the total number of particles).

That the interaction energy is purely local is a significant simplification, arising from the fact that the interactions are short ranged. Minimization of the functional

then poses a purely local problem. At each position, the density $n(\mathbf{r})$ is that which minimizes

$$e[\nu(\mathbf{r})] - \mu_L n(\mathbf{r}) \quad (4)$$

where the local chemical potential is

$$\mu_L(\mathbf{r}) \equiv \mu - V(\mathbf{r}) . \quad (5)$$

For a vortex lattice (at large filling factor), the energy density is

$$e[\nu] = n_0 b \nu^2 \times g_{\text{qH}} , \quad (6)$$

where $b = 1.1596$ is the Abrikosov parameter for a triangular lattice. In this case, the dependence of density on local chemical potential is simple:

$$n(r) = n_0 \mu_L(r)/(2b) , \quad (7)$$

leading to a Thomas-Fermi profile [17] even in the LLL [18].

For filling factors below a critical value ν_c (with ν_c on the order of 10 [3, 19]) the vortex lattice is replaced by groundstates that include incompressible quantum Hall fluids. The energy density is then a complicated function of ν , containing cusps at the incompressible groundstates. The presence of these cusps gives rise to a step-like dependence of n on μ_L . Taken with Eq. (5), this step-like dependence of n on μ_L becomes a step-like dependence in space for a confined system (for which $V(\mathbf{r})$ is not constant in space).

We remark that non-interacting fermions in a rapid rotation regime, and subject to a slowly varying potential, will display density profiles similar to the ones we discuss here [20]. In that case, the result follows from the Landau level structure for non-interacting fermions, while all results discussed in this paper are due to interaction effects. The analogy between the two cases is strengthened by the ‘composite fermion’ formulation of some of our bosonic liquids. This analogy remains incomplete, however, as some of the features that we see (such as the competition between composite fermion states and the paired Moore-Read state) cannot be captured by a model of non-interacting composite fermions.

C. Example: two liquids

As a simple example, we consider the possibility of two competing incompressible states: $\nu_1 = 1/2$ (the Laughlin state, for which the interaction energy for contact interactions vanishes, $\epsilon_1 = 0$ [1]), and $\nu_2 = 2/3$ for which the interaction energy per particle is ϵ_2 . These two liquids are the $p = 1, 2$ members of a series at $\nu_p = p/(p+1)$, which can be understood as integer quantum Hall liquids of composite fermions each consisting of a boson with a single vortex attached [2, 4]. Since the system is bounded, we must also consider the case of no particles, $\nu = 0$. The energy per unit area takes the values

$e[\nu = 0] = e[\nu = 1/2] = 0$, and $e[\nu = 2/3] = (2/3)n_0\epsilon_2$. Minimizing Eq. (4) with respect to these three possible values of the density, $n = \nu n_0$, we find:

$$\begin{aligned} \nu &= 0 & (\mu < 0) \\ \nu &= 1/2 & (0 < \mu < \mu_c) \\ \nu &= 2/3 & (\mu_c < \mu) . \end{aligned} \quad (8)$$

with $\mu_c = 4\epsilon_2$.

Let us take a harmonic confining potential $V(r) = \frac{1}{2}k_2 r^2$, such that $\mu_L(r) = \mu - \frac{1}{2}k_2 r^2$. The $\nu = 1/2$ state forms a disc that extends out to $\mu_L = 0$ (beyond this $\mu_L < 0$ so $\nu = 0$), to a radius $r_1 = \sqrt{2\mu/k_2}$. We further see that there is a critical value of chemical potential, $\mu_c = 4\epsilon_2$, at which the $\nu = 2/3$ state will first appear in the center of the trap (where μ_L is maximum). Evaluating the total number of atoms in the $\nu = 1/2$ disc at μ_c , we find the critical number

$$N_c = (1/2)n_0\pi r_1^2 = 2\epsilon_2/\lambda_2 \quad (9)$$

with $\lambda_2 = \frac{1}{2}k_2\ell_\perp^2$. Above this critical value, the disc of $\nu = 2/3$ has a radius $r_2 = \sqrt{2(\mu - \mu_c)/k_2}$.

Thus, as the number of particles increases, at first they will form a disc of uniform density $1/2 n_0$ the radius of which increases with increasing N (μ). Once N reaches N_c (or μ reaches μ_c) a second, inner, disc of density $2/3 n_0$ will form, and start to expand with increasing N (or μ). In terms of N and N_c the two steps for $N > N_c$ are located at radii

$$r_2/\ell_\perp = \sqrt{\frac{3}{2}(N - N_c)} , \quad r_1/\ell_\perp = \sqrt{\frac{3}{2}N + \frac{1}{2}N_c} . \quad (10)$$

The energy as a function of N takes the form

$$E(N) = \lambda_2 \left(N^2 - \frac{1}{4}\Theta(N - N_c)(N - N_c)^2 \right) \quad (11)$$

We note that the second derivative of $E(N)$ is discontinuous at $N = N_c$.

Note that, although we have only considered the possibilities that $\nu = 0, 1/2, 2/3$ (and we have not allowed for intermediate values), we do not expect intermediate densities to appear in the density distribution. Since $e[\nu] = 0$ for $\nu < 1/2$, values of ν between 0 and $1/2$ are only possible precisely at $\mu_L = 0$ and hence have no finite extent in a generic inhomogeneous system. For the range, $1/2 < \nu < 2/3$, numerical studies indicate that $e[\nu]$ lies above the straight line joining the end-points $e[\nu = 1/2] = 0$ and $e[\nu = 2/3] = 2/3 n_0\epsilon_2$. Therefore, intermediate filling factors are unstable to phase separation into $\nu = 1/2$ and $\nu = 2/3$.

A slightly different perspective on the formation of two spatially separated liquids is the following. If one imagines adding more and more particles to a $\nu = 1/2$ Laughlin droplet in a (radially symmetric) potential, one will eventually reach the point where the energy of an additional particle at the edge of the Laughlin droplet exceeds

the energy gap Δ_1 to create a single particle in the bulk Laughlin liquid at the center of the droplet. The corresponding critical value for N is $\tilde{N}_c = \Delta_1/(2\lambda_2)$. Using numerical results for Δ_1 and ϵ_2 (as given, for example in [4]) $\tilde{N}_c$ is found to be slightly higher than N_c , in agreement with the fact that a $\nu = 1/2$ system with additional quasi-particles will lower its energy by phase separating into a central $\nu = 2/3$ liquid and an outer ring of $\nu = 1/2$ liquid.

Repeating the analysis of the co-existing $\nu = 1/2$ and $\nu = 2/3$ liquids in a quartic potential $V_4(r) = \frac{1}{4}k_4 r^4$ leads to a critical N of

$$N_c = \sqrt{\frac{\epsilon_2}{\lambda_4}}, \quad (12)$$

with $\lambda_4 = \frac{1}{4}k_4\ell_\perp^2$, and with inner and outer edges at

$$\begin{aligned} (r_2/\ell_\perp)^2 &= \frac{3}{4}(\sqrt{9N^2 - 8N_c^2} - N), \\ (r_1/\ell_\perp)^2 &= \frac{9}{4}N - \frac{1}{4}\sqrt{9N^2 - 8N_c^2}. \end{aligned} \quad (13)$$

The condition that $V(r)$ be slowly varying, phrased as $\frac{dV(r)}{dr}\ell_\perp \ll g_{\text{QH}}$, is equivalent to demanding that $r_i/\ell_\perp \gg 1$ for all i . For this to be possible, a total number of particles on the order of $N = 100$ or more is required. This puts the quantized density profiles out of the reach of exact diagonalization studies.

D. More liquids

Including in the analysis more quantum liquids, at filling factors $\nu_1 < \nu_2 < \dots$ leads to a sequence of critical values $N = N_c^{(k)}$, marking the start of the formation of a central region of a quantum Hall fluid at $\nu = \nu_k$.

To illustrate the step-like profiles for higher filling factors, we use the interaction energy function $e[\nu]$ calculated by exact diagonalization studies on a torus [3]. In the graphs presented in Fig. 1, a confining potential $V(r) = \frac{1}{2}k_2 r^2$ is assumed. The energy functional used is for system size of $N_V = 6$ single-particle states on a torus [for which the numerical results extend up to $\nu = 25/3$].

For filling factors larger than $\nu_c \simeq 6$, the groundstate is a compressible vortex lattice [3]. The steps seen in the density distribution in that regime are an artifact of the minimum change of filling factor, $\Delta\nu = 1/6$, allowed by the numerics on a finite system with $N_V = 6$. In this regime, the density profile will be smooth (we emphasize again that our results apply only on scales large compared to $\ell_\perp$, so do not describe the short-range density modulations associated with the unit cell of a triangular vortex lattice).

For filling factors less than $\nu_c \sim 6$, these calculations show plateaus appearing at the densities of a set of incompressible quantum Hall fluids, including the sequence of Moore-Read and Read-Rezayi states at $\nu = k/2$




FIG. 1: Density profiles in a single layer, at subcritical rotation, parametrized by density at the center of the trap. The results are for an energy functional obtained from exact diagonalizations of a system with $N_V = 6$ states in toroidal geometry (aspect ratio $a/b = \sqrt{3}/2$).




FIG. 2: Density profiles in a single layer, for cases with maximum central densities of up to $\nu = 1$. The energy functional is for $N_V = 12$ states in toroidal geometry (aspect ratio $a/b = 1$). The quantum liquid at $\nu = 1$ is the Moore-Read paired state and the $\nu = 1/2$ state is the Laughlin liquid. In between, a number of composite fermion states (at $\nu = 2/3, 3/4$) are observed.

(black dashed lines) as well as some of the CF sequence $\nu = p/(p+1)$ (black dotted lines). We note that finite size effects in these results for $N_V = 6$ can affect the widths and positions of the plateaus appearing in Fig. 1. This figure illustrates the general feature of the formation of plateaus at the densities of incompressible states. One interesting detail is the competition between successive members of the CF series at $\nu = p/(p+1)$, and the Moore-Read (MR) liquid[3, 4, 21] at $\nu = 1$. These states have very different correlations, and one does not expect the higher CF states to be stable against the pairing in-

stability that is at the basis of the formation of the MR state. This competition is highlighted in Fig. 2, which shows the density profiles for systems with maximum central densities of up to $\nu = 1$ calculated for a system with $N_V = 12$ vortices (which, unlike $N_V = 6$, also permits the appearance of $\nu = 3/4$, and for which finite size effects on the plateau widths are much reduced). Note that there is no appearance of an incompressible state at $\nu = 5/6$. This indicates that, at least within this small system calculation, the $\nu = 5/6$ state appears to be unstable to phase separation into the $\nu = 3/4$ and $\nu = 1$ states.

III. MULTI-LAYER GEOMETRY

A. Experimental set-up

We now investigate the situation where an optical lattice in the z direction is imposed on a cigar shaped cloud of atoms rotating around the z -axis. If the optical potential is sufficiently deep, this will define a stack of parallel planes with (weak) tunneling t between the planes. The idea is that parameters can be chosen such that, while the original cloud is in a mean field regime with total filling factor $\nu > \nu_c$ such that it would display a vortex lattice, the individual layers defined by the optical lattice can be in a quantum regime, so that the entire configuration becomes a stack of weakly coupled quantum liquids. We will assume the presence of a chemical potential $\mu(z) = \mu_0 - \mu_2(z/d)^2$ (with d the distance between the layers), which will induce a slow modulation in the number of particles per layer. One may view a set-up like this as a road towards isolating a single layer in the quantum Hall regime, or to perform diagnostics on the entire multi-layer geometry.

Reaching the regime where quantum Hall states can form requires a number of conditions. First of all, to avoid the formation of a mean-field vortex lattice, the filling factor in each of the layers should be sufficiently low (meaning high-enough total angular momentum per particle) and the interlayer coupling t should be sufficiently small. The second condition is that the tunneling is well below the gap expected for the quantum Hall states.

Concerning the first condition, a quick estimate is made as follows. The critical filling factor ν_c for a single layer has been found to be of the order $\nu_c \sim 10$. Assuming N particles, N_V vortices and N_L layers, quantum melting will be possible if $N/(N_L N_V) < \nu_c$. Taking, for example, $N = 5000$ particles and $N_V = 100$ vortices (filling factor $\nu = 50$), slicing up the condensate into, say, $N_L = 50$ layers of 100 particles each would produce a filling factor per layer ν' of order unity. For this choice of parameters, a gradual lowering of the tunneling amplitude causes a quantum melting of the vortex lattice at a critical $t = t_{c1}$. For $t < t_{c1}$ one expects a state with quantum liquids in each of the layers.

The quantum Hall gap that features in the second con-

dition is set by the interactions between the atoms (as encoded in the scattering length a_s) and the length scales $\ell_\perp$ and $\ell_\parallel$ of the confinement in the $x - y$ and z directions. (In the multi-layer case, $\ell_\parallel$ refers to the thickness of a single layer.) A theoretical estimate[3, 4] gives

$$\Delta \simeq 0.1 \, 4\pi g_{\text{qH}} \simeq \frac{a_s}{\ell_\parallel} \hbar \omega_\perp. \quad (14)$$

with g_{qH} as in Eq. (2). With the parameters as given below Eq. (2), the quantum Hall gap is found to be in the order of 1 nK.

There are several possibilities for experimental detection of the formation of incompressible states in the multi-layer set-up. One possibility, which we worked out in [16], is to focus on the total particle number per layer, *i.e.* on the density profile as a function of the layer coordinate $\rho(z)$. In this paper we focus on a detection scheme where one allows a rapid expansion of the system and then measures the density profile as a function of the radial coordinate r .

B. Quantum melting of the vortex lattice

We now turn to the analysis of the quantum melting of the vortex lattice in the multi-layer geometry. For this, we will assume that all layers have identical axial and radial confinement, with a chemical potential independent of z . If the filling factor per layer, $\nu' = N/(N_L N_V)$, is well below the critical filling factor ν_c for a single layer, isolated single layers will be in the regime where the vortex lattice has melted. This means that for this situation, a gradual lowering of t (by turning up the strength of the optical lattice potential), will lead to a melting of the vortex lattice at a critical tunneling $t = t_{c1}$. Equivalently, there is a critical layer filling factor $\nu'_c(t)$ for given tunneling t . We will use two inequivalent techniques to find the value of $\nu'_c(t)$.

Due to the short-range nature of the interaction, the inter-layer interactions are much smaller than both the intra-layer interaction and the tunneling, mean-field theory then predicts that the vortex lattices in different layers will lie on top of each other. Note that this is very different from the quantum Hall multi-layer setting in semiconductor devices, where the long-ranged Coulomb interaction can dominate the tunneling.

We model the coupled layers by including a Josephson term:

$$H = \sum_i H_i^{(0)}(\mu) - \sum_{\langle ij \rangle} \int d^2 \mathbf{r} \, t (\psi_i^\dagger \psi_j + \psi_j^\dagger \psi_i), \quad (15)$$

where we include tunneling only between neighboring layers, labeled by i .

Our first approach for determining $\nu'_c(t)$ is to extend the single layer effective theory of vortex fluctuations derived by Sinova *et al.* [19] to the multi-layer case. We




FIG. 3: Critical layer filling factor $\nu'_c(t)$ as a function of $\ln(t/(\nu'g_{qH}))$, obtained by applying a Lindemann criterion to the vortex fluctuations (Eq. 17) in the limit $N_L \rightarrow \infty$, $N_V \rightarrow \infty$.

neglect the gapped sound mode[22] and focus on the low-energy Tkachenko modes. In terms of the deviations $\mathbf{u}^i(\mathbf{r})$ of the vortex positions from their equilibrium values, the Josephson coupling takes the following form:

$$\int_0^\infty d\tau \int d\mathbf{q} \sum_{\langle ij \rangle} \left[\frac{1}{2} t \frac{\alpha_1}{q^2} \left(|u_L^i - u_L^j|^2 + |u_T^i - u_T^j|^2 \right) \right], \quad (16)$$

with an additional (constant) contribution to the chemical potential. In addition to the vortex positions, each layer has a phase φ_i . We assume that the phases of all layers are identical, an approximation which is valid when $t \gg t_{c2}$.

On distances smaller than the axial healing length (in units of the optical lattice period d) $\xi_{\parallel} \approx \sqrt{t/g_{qH}\nu'}$, all bosons are in the same axial state. Therefore, they behave as a single layer and the vortices are ‘rigid’ on the scale of $\xi_{\parallel}$. In order to capture this effect, we use $\xi_{\parallel}$ as a short-distance cut-off in the axial direction. The mean square displacement of a vortex is given by:

$$\langle u^2 \rangle \approx \frac{1}{2N_L} \sum_k \int \frac{d^2\mathbf{q}}{(2\pi)^2} \frac{q^2}{\alpha_1} \left(\sqrt{\frac{K_L}{K_T}} + \sqrt{\frac{K_T}{K_L}} \right) e^{-|k|\xi_{\parallel}}, \quad (17)$$

where N_L is the number of layers, $k = 0, 2\pi/N_L, \dots$ is expressed in units of the inverse layer spacing, $1/d$, and q is in units of $1/\ell_{\perp}$. The constants α_1 , α_2 and c_{66} are as in Ref. [19] and

$$K_L(\mathbf{q}, k) = \frac{\alpha_2}{q^2} + \frac{1}{2} \frac{\alpha_1}{q^2} t (1 - \cos(k)) \quad (18)$$

$$K_T(\mathbf{q}, k) = c_{66} q^2 + \frac{1}{2} \frac{\alpha_1}{q^2} t (1 - \cos(k)). \quad (19)$$

The Lindemann criterion predicts the quantum melting transition to occur when the mean square displacement




FIG. 4: Critical filling factor per layer $\nu'_c(t)$ as a function of $\ln(t/(\nu'g_{qH}))$ for $N_L = 8, 50$ layers and $N_V = 10, 100$ vortices. The solid line indicates the limit $N_L \rightarrow \infty$, $N_V \rightarrow \infty$.

is[23]

$$\langle u^2 \rangle \approx \alpha_L^2 a_0^2, \quad (20)$$

where $\alpha_L \approx 0.1 - 0.2$ is a system-dependent (*a priori* unknown) constant and a_0 is the lattice spacing. We take $\alpha_L = 0.15$ and have plotted the resulting critical layer filling factor $\nu'_c(t)$ in Fig. 3.

Our second approach to estimate the critical filling factor per layer $\nu'_c(t)$ is based on the depletion of the condensate [19]. This criterion is, perhaps, less direct in determining the melting point of the vortex lattice than the Lindemann criterion, and may tend to overestimate the critical filling factor $\nu'_c(t)$. However, given its simplicity, it is valuable to know the results of the calculation based on this criterion.

The filling factor $\tilde{\nu}$ for bosons outside the condensate is:

$$\tilde{\nu} - \nu = \frac{\ell_{\perp}^2}{4\pi} \frac{1}{N_L} \sum_k \int_{BZ} d\mathbf{q} (\xi_{k,\mathbf{q}}/E_{k,\mathbf{q}} - 1), \quad (21)$$

The one-layer expressions [19] ($\xi_{\mathbf{q}}^0$, $\lambda_{\mathbf{q}}^0$) are generalized by $\xi_{k,\mathbf{q}} = \xi_{\mathbf{q}}^0 + t(1 - \cos(k))$, $E_{k,\mathbf{q}} = \sqrt{\xi_{k,\mathbf{q}}^2 - |\lambda_{\mathbf{q}}^0|^2}$. The $k = 0$ integrand diverges when $q \rightarrow 0$, due to phase fluctuations. In a finite system we can use the system size as an infrared cutoff, $q_{min} = 1/\sqrt{2\pi N_V}$. We equate the quantum melting point with the filling factor where half the bosons are in an excited state and find the critical filling factors shown in Fig. 4.

The critical filling factor for large tunneling, determined in this way, is well approximated by $\nu'_c(t \rightarrow \infty) = \nu'_c(t \rightarrow 0)/N_L$, in agreement with the intuition that for large tunneling the system behaves as a single layer with total filling factor $\nu = N_L \nu'$. As can be observed in Fig. 4, in a large range of tunneling amplitudes the critical filling factor is independent of N_L and N_V . Eq. (21)

can be approximated by a converging integral, which is exact for $N_L \rightarrow \infty$, $N_V \rightarrow \infty$. For the parameters quoted above, $N_L = 50$, $N_V = 100$, $\nu' = 1$, we find that the critical tunneling for quantum melting of the vortex lattice is $t_{c1} \approx 0.18g_{\text{qH}}$.

C. Quantum phases of the multi-layer systems

The melting of the vortex lattice leaves a multi-layer system of highly correlated quantum liquids. The nature of the quantum liquids is not *a priori* clear. In the limit of vanishing tunneling, the isolated layers will display single-layer behavior. The states formed close to the melting transition are, however, not necessarily adiabatically connected to these but could be separated by a quantum phase transition.

For t smaller than but rather close to t_{c1} there is the possibility for correlated multi-layer states. For the case of electronic quantum Hall systems, such correlated multi-layer states have been proposed and analyzed (an early reference is [24]). For the atomic systems the multi-layer states can be very different, as there are no interlayer interactions and interlayer correlations are driven by tunneling alone.

Our main interest is in the regime where the individual layers are weakly coupled. In this regime, the atoms will be able to move in between layers and form an equilibrium density profile. In this regime we expect a local density approximation in both the coordinates z and r to be valid. In section III D below, we analyze this regime and work out the density profile arising after expansion.

For $t \ll g_{\text{qH}}$, interlayer transport will stop and individual layers can be at a sharply quantized particle number, realizing what we call a quantum Hall-Mott (qH-Mott) phase. To determine the boundaries of these qH-Mott phases, we have analyzed a simple effective Bose-Hubbard model. In the simplest case we take quadratic in-plane confinement and assume that all layers form a Laughlin state (with filling factor per layer $\nu' = 1/2$). For this situation we obtained the following expression for the critical tunneling t for entering the qH-Mott state with M particles,

$$t_{c2}(\mu) = -\lambda_2 \left[M + \frac{1}{2} - \frac{\mu}{2\lambda_2} \right] \left[M - \frac{1}{2} - \frac{\mu}{2\lambda_2} \right], \quad (22)$$

for a chemical potential in the interval $M - \frac{1}{2} \leq \frac{\mu}{2\lambda_2} \leq M + \frac{1}{2}$. As in the standard Bose-Hubbard model, the curves $t_c(\mu)$ define characteristic ‘lobes’ in the μ - t plane, marking the boundaries of the qH-Mott phases. We observe that $t_{c2}(\mu)$ has a maximum of $\lambda_2/4$, which is of order g_{qH}/M . This scale is in agreement with the fact that the lowest excitations of the quantum Hall droplets are edge excitations with energies of order g_{qH}/M .

In Fig. 5 we display $\mu(z)$ - t diagrams for Laughlin states in quadratic and quartic in-plane confinement. In the quartic case, the maximum of t_{c2} grows with μ . This




FIG. 5: Phase diagrams in the $\mu(z)$ - t plane, for a situation with $\nu = 1/2$ Laughlin quantum Hall states in the layers. The shaded areas correspond to quantum Hall Mott states, where the particle number per layer is sharply quantized. In the remaining part of the diagram there are interlayer fluctuations, giving rise to an axial superfluid. The top and bottom panels are for quadratic and quartic confinement, respectively.

makes possible a situation where the central layers of the stack (near $z = 0$) display qH-Mott states, while the outer layers are in the fluctuating regime.

We wish to emphasize that these quantum Hall-Mott phases are related to the well known Mott phase. When a 1D lattice of atoms in the Mott phase is subjected to rotation (with fixed M), each site will expand to a layer with a well defined filling factor ν' . When $\nu' \approx \nu_c$, the single layer critical filling factor, the system will undergo a quantum phase transition to the quantum Hall-Mott phase. This transition occurs when $t < t_{c2}$, where we extend the definition of t_{c2} to the Mott phase.

An interesting point in the $t - \nu'$ phase diagram (for fixed M) is $(t, \nu') = (t^*, \nu^*)$, where $t_{c1} = t_{c2} \equiv t^*$ and ν^* is the critical filling factor for $t = t^*$. At this point, the Mott transition coincides with the quantum melting transition.

D. Density profile after expansion

In this section we focus on the regime where the tunneling is such that we have a stack of weakly coupled quantum Hall layers. We can apply the ideas developed in section II, where we showed how each individual layer is built from local pieces of quantum Hall fluid. Having a slowly varying chemical potential $\mu(z)$ implies that the density landscape varies slowly from layer to layer,

and that processes where atoms tunnel from the edge of a local quantum Hall fluid in one layer i to the corresponding edge in an adjacent layer $i \pm 1$, are possible. The energy scale for such inter-layer edge excitations is of order g_{qH}/N_i , with N_i the number of particles in layer i . We assume that the tunneling t is large on this scale, such that tunneling events can establish an equilibrium density profile $n(r, z)$.

The density profile can be written as

$$n(r, z) = n_0 \nu [\mu_0 - \mu_2 (z/d)^2 - V(r)] , \quad (23)$$

with $\nu[\mu] = \nu_i$ for $\mu_c^{(i)} < \mu < \mu_c^{(i+1)}$. A measurement of particle number per layer averages this over r :

$$\rho(z) = \int 2\pi dr n(r, z) . \quad (24)$$

The profiles $\rho(z)$ have been discussed in [16]. Here we focus instead on the density profile arising after switching off the optical lattice potential in the z -direction and the trapping potential. In this expansion, each tightly confined layer expands enormously in the z direction, until all layers overlap. The expansion image is thus obtained by averaging over z ,

$$\rho(r) = \int dz n(r, z) . \quad (25)$$

Expansion in the radial direction, which is needed for an actual measurement of $\rho(r)$, will not affect the density profile other than by an overall expanding scale[13].

The integral in Eq. 25 is easily worked out. If we assume quadratic in-plane confinement, $V(r) = \frac{1}{2}k_2 r^2$, the profile takes the form of a sum of semi-circle arcs

$$\rho(r) = n_0 \left(\frac{2k_2}{\mu_2} \right)^{1/2} \sum_i [\nu_i - \nu_{i-1}] \sqrt{r_i^2 - r^2} \quad (26)$$

with

$$r_i = \left(\frac{2(\mu_0 - \mu_c^{(i)})}{k_2} \right)^{1/2} . \quad (27)$$

For quartic confinement the result is

$$\rho(r) = n_0 \left(\frac{k_4}{\mu_2} \right)^{1/2} \sum_i [\nu_i - \nu_{i-1}] \sqrt{r_i^4 - r^4} \quad (28)$$

with

$$r_i = \left(\frac{4(\mu_0 - \mu_c^{(i)})}{k_4} \right)^{1/4} . \quad (29)$$

It is clear that, while the expansion does wash out the sharp steps in the density profiles, it leaves cusps in the averaged profiles. For in-plane confinement $V(r) \propto r^\alpha$ these cusps take the form of square root singularities, $\rho(r) \propto \sqrt{R^\alpha - r^\alpha}$. In Figs. 6 and 7 we display some of




FIG. 6: Radial density profile $\rho(r)$, for a system with quantum Hall liquids at $\nu_1 = 1/2$, $\nu_2 = 2/3$, $\nu_3 = 3/4$ and $\nu_4 = 1$. These profiles arise after expansion in the z -direction. The top panel is for quadratic in-plane confinement, with parameters chosen such that the landscape in the central layer (at $z = 0$) coincides with the top curve in Fig. 2. The bottom panel assumes quartic confinement.

the profiles $\rho(r)$. The positions of the cusps provide information on the incompressible liquid states. Fitting the measured density profile to (26), involving the set of filling factors ν_i , allows the critical chemical potentials, μ_c^i , to be measured. This would provide direct information on the energetics and incompressibility of the correlated quantum liquids.

We now estimate the magnitude of fluctuations that affect these mean field density profiles. Focusing on a qH fluid (ν_k) in one of the layers, we can evaluate the energy for a displacement of the outer edge, $r_k \rightarrow r_k + \delta r_k$. In evaluating this energy, we assume that the chemical potential at the center of the layer is of order g_{qH} (which is the natural scale in the present regime). With that we find

$$\delta E \simeq g_{\text{qH}} \left(\frac{\delta r_k}{\ell_\perp} \right)^2 \quad (30)$$

Equating this to the scale set by the tunneling gives the estimate

$$\delta r_k \simeq \ell_\perp \sqrt{\frac{t}{g_{\text{qH}}}} . \quad (31)$$




FIG. 7: Radial density profile $\rho(r)$, for a system with quantum Hall liquids at $\nu_1 = 1/2$ and $\nu_2 = 2/3$, for quartic in-plane confinement.

We conclude that, for t of order g_{qH} or smaller, the fluc-

tuations induced on r_k are of order $\ell_\perp$ or less. This means that, in the presence of tunneling, the steps within each of the layers remain well-defined on a scale $\ell_\perp$. Accordingly, one expects that the cusps in the profile after expansion will be recognizable if the overall extent (in the x - y plane) of the original cloud is large on the scale of $\ell_\perp$.

The features in $\rho(r)$ can be sharpened further by eliminating some of the outer layers from the stack, keeping only a number of layers centered around $z = 0$, or by having an axial confinement $\mu(z)$ that is steeper than quadratic.

We thank J. Dalibard and V. Schweikhard for illuminating discussions on the prospects for experimental realization of atomic quantum Hall states. This research was supported by the Netherlands Organisation for Scientific Research, NWO, the Foundation FOM of the Netherlands (JWR, FJMvL and KS), and by EPSRC grant GR/S61263/01 (NRC).

-
- [1] N.K. Wilkin, J.M.F. Gunn and R.A. Smith, Phys. Rev. Lett. **80**, 2265 (1998); N.K. Wilkin and J.M.F. Gunn, Phys. Rev. Lett. **84**, 6 (2000).
 - [2] N.R. Cooper and N.K. Wilkin, Phys. Rev. **60**, R16279 (1999).
 - [3] N.R. Cooper, N.K. Wilkin, and J.M.F. Gunn, Phys. Rev. Lett. **87**, 120405 (2001).
 - [4] N. Regnault and Th. Jolicoeur, Phys. Rev. Lett. **91**, 030402 (2003); Phys. Rev. B **70**, 241307(R) (2004); Mod. Phys. Lett. B **18**, 1003 (2004); C.-C. Chang, N. Regnault, Th. Jolicoeur and J.K. Jain, cond-mat/0412253.
 - [5] T.-L. Ho and E.J. Mueller, Phys. Rev. Lett. **89**, 050401 (2002).
 - [6] B. Paredes, P. Zoller and J.I. Cirac, Phys. Rev. A **66**, 033609 (2002).
 - [7] J.W. Reijnders, F.J.M. van Lankvelt, K. Schoutens and N. Read, Phys. Rev. Lett. **89**, 120401 (2002); Phys. Rev. A **69**, 023612 (2004).
 - [8] V. Bretin, S. Stock, Y. Seurin and J. Dalibard, Phys. Rev. Lett. **92**, 050403 (2004).
 - [9] V. Schweikhard, I. Coddington, P. Engels, V.P. Mogen-dorff and E.A. Cornell, Phys. Rev. Lett. **92**, 040404 (2004).
 - [10] B. Paredes, P. Fedichev, J.I. Cirac and P. Zoller, Phys. Rev. Lett. **87**, 010402 (2001).
 - [11] J. Sinova, C.B. Hanna and A.H. MacDonald, Phys. Rev. Lett. **90**, 120401 (2003).
 - [12] M.A. Cazalilla, Phys. Rev. A **67**, 063613 (2003); M.A. Cazalilla, N. Barberán and N.R. Cooper, Phys. Rev. B **71**, 121303(R) (2005).
 - [13] N. Read and N.R. Cooper, Phys. Rev. A **68**, 035601 (2003); E. Altman, E. Demler and M.D. Lukin, Phys. Rev. A **70**, 013603 (2004).
 - [14] F.J.M. van Lankvelt, J.W. Reijnders and K. Schoutens, presentation S3.003 at 2004 APS March meeting (Montreal).
 - [15] N.R. Cooper, presentation at the conference ‘Quantum Gases’ KITP Santa Barbara (May 2004).
 - [16] N.R. Cooper, F.J.M. van Lankvelt, J.W. Reijnders and K. Schoutens, *Density profiles for atomic quantum Hall states*, cond-mat/0409146 (unpublished).
 - [17] A.L. Fetter, Phys. Rev. A **64**, 063608 (2001); U.R. Fischer, G. Baym, Phys. Rev. Lett. **90**, 140402 (2003).
 - [18] G. Watanabe, G. Baym and C.J. Pethick, Phys. Rev. Lett. **93**, 190401 (2004); N.R. Cooper, S. Komineas and N. Read, Phys. Rev. A **70**, 033604 (2004).
 - [19] J. Sinova, C.B. Hanna and A.H. MacDonald, Phys. Rev. Lett. **89**, 030403 (2002).
 - [20] T.-L. Ho and C.V. Ciobanu, Phys. Rev. Lett. **85**, 4648 (2000).
 - [21] G. Moore and N. Read, Nucl. Phys. B **360**, 362 (1991).
 - [22] S.A. Gifford and G. Baym, Phys. Rev. A **70**, 033602 (2004).
 - [23] A. Rozhkov and D. Stroud, Phys. Rev. B **54**, R12697 (1996).
 - [24] X. Qiu, R. Joynt and A.H. MacDonald, Phys. Rev. B **40**, 11943 (1989).